

Supplementary Table 1. Characteristics of patients for whom VATS sleeve lobectomy was attempted

Characteristic	Intention to treat VATS (n=19)
Age (yr)	59.7±14
Male sex	17 (89.5)
Smoking history	
Never smoker	3 (15.8)
Ex-smoker	8 (42.1)
Current smoker	8 (42.1)
Pulmonary function test	
FEV1 (%PRED)	79.5±15.9
DLCO (%PRED)	81.7±16.1
Comorbidities	
Diabetes	3 (15.8)
Hypertension	6 (31.6)
Chronic lung disease	5 (26.3)
Cerebrovascular disease	2 (10.5)
Chronic liver disease	3 (15.8)
Previous malignancy	1 (5.3)
CCI score	
0	7 (36.8)
1	9 (47.4)
2	0 (0.0)
≥3	3 (15.8)
Tumor location	
RUL	4 (21.1)
LUL	12 (63.2)
LLL	3 (15.8)
Clinical T stage	
T1	3 (15.8)
T2	15 (78.9)
T3	1 (5.3)
Clinical N stage	
N0	11 (57.9)
N1	7 (36.8)
N2	1 (5.3)

Values are presented as mean±standard deviation or number (%). VATS, video-assisted thoracoscopic surgery; FEV1, forced expiratory volume in 1 second; PRED, predicted; DLCO, diffusing capacity for carbon monoxide; CCI, Charlson comorbidity index; RUL, right upper lobe; LUL, left upper lobe; LLL, left lower lobe.