

**Supplementary Table 2.** Perioperative outcomes of patients for whom VATS sleeve lobectomy was attempted

Variable	Intention to treat VATS (n=19)
Histology	
Squamous cell carcinoma	15 (78.9)
Mucoepidermoid carcinoma	2 (10.5)
Atypical carcinoid tumor	1 (5.3)
Extranodal marginal zone B cell lymphoma	1 (5.3)
Tumor size (cm)	3.3±1.5
Pathologic T stage	
T1	6 (31.6)
T2	12 (63.2)
T3	1 (5.3)
Pathologic N stage	
N0	11 (57.9)
N1	6 (31.6)
N2	2 (10.5)
Lymphatic invasion	7 (36.8)
Vascular invasion	1 (5.3)
Perineural invasion	5 (26.3)
Adjuvant therapy	
Concurrent chemoradiation	1 (5.3)
Chemotherapy	7 (36.8)
R0 resection	19 (100)
Bronchial margins (cm)	1.1±1.1
Total resected lymph nodes	19.7±8.2
Positive lymph nodes	0.8±1.2
Operating time (min)	246±69
Intensive care unit stay (day)	1.7±1.9
Chest tube duration (day)	6.1±2.1
Postoperative hospital stay (day)	8.9 (8–10)

Values are presented as number (%), mean ± standard deviation, or median (interquartile range).

VATS, video-assisted thoracoscopic surgery.